



Downloaded from: <http://bucks.collections.crest.ac.uk/>

This document is protected by copyright. It is published with permission and all rights are reserved.

Usage of any items from Buckinghamshire New University's institutional repository must follow the usage guidelines.

Any item and its associated metadata held in the institutional repository is subject to

Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0)

Please note that you must also do the following;

- the authors, title and full bibliographic details of the item are cited clearly when any part of the work is referred to verbally or in the written form
- a hyperlink/URL to the original Insight record of that item is included in any citations of the work
- the content is not changed in any way
- all files required for usage of the item are kept together with the main item file.

You may not

- sell any part of an item
- refer to any part of an item without citation
- amend any item or contextualise it in a way that will impugn the creator's reputation
- remove or alter the copyright statement on an item.

If you need further guidance contact the Research Enterprise and Development Unit
ResearchUnit@bucks.ac.uk

Manuscript has been accepted but not published:
Accepted author manuscript version reprinted, by permission, from
International Journal of Sports Physiology and Performance (IJSP), 2020
(ahead of print). © Human Kinetics, Inc.

Manuscript has been accepted but not published:

Accepted author manuscript version reprinted, by permission, from Journal of Sports
Physiology and Performance, 2020 (ahead of print). © Human Kinetics, Inc.

Abstract

Purpose: The aim of this study was to investigate the relationship between selected anthropometric, physiological, and upper body strength measures and 15-km handcycling time trial performance.

Methods:

Thirteen, trained H3/H4 male handcyclists performed a 15-km time trial, graded exercise test, 15-s all-out sprint and one repetition maximum assessment of bench press and prone bench pull strength. Relationship between all variables were assessed using a Pearson's correlation coefficient matrix with mean time trial velocity representing the principal performance outcome.

Results:

Power at a fixed blood lactate concentration of $4 \text{ mmol}\cdot\text{l}^{-1}$ ($r = .927$; $p < 0.01$) showed an extremely large correlation to TT performance, whilst relative $\text{VO}_{2\text{peak}}$ ($r = .879$; $p < 0.01$), power-to-mass ratio ($r = .879$; $p < 0.01$), peak aerobic power ($r = .851$; $p < 0.01$), gross mechanical efficiency ($r = .733$; $p < 0.01$), relative prone bench pull strength ($r = .770$; $p = 0.03$) relative bench press strength ($r = .703$, $p = 0.11$), and maximum anaerobic power ($r = .678$; $p = 0.15$) all demonstrated a very large correlation with performance outcomes.

Conclusion: Findings of the present study indicate that power at a fixed blood lactate concentration of $4 \text{ mmol}\cdot\text{l}^{-1}$, relative $\text{VO}_{2\text{peak}}$, power-to-mass ratio, peak aerobic power, gross mechanical efficiency, relative upper body strength, and maximum anaerobic power are all significant determinants of 15-km TT performance in H3/H4 handcyclists.

27 **Introduction**

28 Since its formal recognition as a sport by the International Paralympic Committee in 1999, the
29 popularity of handcycling as both a recreational and competitive sport has grown substantially.³ The
30 sport was first included in the summer 2004 Paralympic Games in Athens and is now also
31 incorporated within the wheelchair classifications of Para-triathlon, a sport that debuted at the
32 summer 2016 Paralympic Games in Rio de Janeiro. Handcyclists are classified into one of five
33 categories (H1 - H5) according to the nature of their physical impairment, with H1 athletes typically
34 having the greatest physical impairment and lowest physical function.³³ Handcycling can be viewed
35 as an endurance sport whereby athletes typically compete in road-races and/or individual time trials
36 (TT) over distances ranging from 37 to 80-km and 10 to 35-km, respectively.³³ Due to varying race
37 formats, terrain, tactics and speeds, handcyclists often use variable pacing strategies, such as frequent
38 short accelerations to push opponents, or drafting behind other riders during road races to reduce
39 overall energy cost.^{1,3} Arguably, a TT represents the most pure challenge to a cyclist as, in this format,
40 athletes ride alone, against the clock. Thus, racing tactics briefly mentioned above become irrelevant
41 and a cyclist's ability is laid bare.

42
43 As with many other Paralympic sports, the performance level of handcycling at the elite level has
44 increased considerably, which underlines the importance of optimising training in order to achieve
45 success.²³ Handcycling performance is ultimately dependent upon the physical capabilities of the
46 individual, the design of the handbike, and the interaction between the rider and their equipment,
47 typically referred to as the handbike-user interface.³¹ Whilst the biomechanics,^{20,27,31} handbike-user
48 interface^{4,5,6,22,29,32,34} and physiological characteristics of handcycling performance, namely peak
49 oxygen uptake ($\dot{V}O_{2\text{peak}}$), peak aerobic power output (PO_{peak}) and gross mechanical efficiency (GME)
50 have been extensively investigated.^{1,9,13,19,21,25} To date, only a handful of studies have examined the
51 relationship between the aforementioned physiological characteristics and handcycling race

performance. Janssen et al,¹⁹ investigated how the physical capacity of sixteen male handcyclists with either spinal cord injury, spina bifida or lower limb amputation related to performance during a 10-km handcycling race. Relative $\dot{V}O_{2peak}$ ($r = .90, p < 0.01$), PO_{peak} ($r = .91, p < 0.01$), and power-to-mass ratio ($r = .89, p < 0.01$) were all demonstrated to be significant predictor of 10-km race performance. Lovell et al,²¹ studied the aerobic performance of ten trained and ten untrained male handcyclists with spinal cord injury. They reported that PO_{peak} ($r = .87, p < 0.01$), $\dot{V}O_{2peak}$ ($r = .67, p = 0.03$), and GME ($r = .50, p = 0.04$) were the best predictors of handcycling performance during a laboratory-based, 20-km TT. More recently, Fischer et al,¹³ examined the physiological determinants of handcycling performance of seven, male, H2 handcyclists and found that $\dot{V}O_{2peak}$ ($r = .89, p < 0.01$), $\dot{V}O_{2VT1}$ ($r = .96, p < 0.01$), $\dot{V}O_{2VT2}$ ($r = .92, p = 0.03$) and PO_{peak} ($r = .85, p = 0.02$) were all significantly correlated with performance during a simulated 22-km TT.

Whilst extensively examined in able-bodied cycling,^{10,11} few studies have investigated the impact of anaerobic threshold, anaerobic capacity and upper-body strength upon handcycling performance. The attainment of high-power output during an all-out sprint is frequently used as a measure of anaerobic performance in able-bodied cycling^{10,11} and Quittman et al,²⁶ recently suggested that an understanding of a handcyclists physiological profile should be augmented by testing maximal anaerobic power ($PO_{max,AO15}$). In cycling, anaerobic capacity can be defined as the difference between $PO_{max,AO15}$ and PO_{peak} commonly known as the anaerobic power reserve (APR). Several authors have demonstrated that decrements observed in all-out cycling performance seems to conform to a general relationship with a single exponential decay model, which describes a decrement in power versus increasing duration.^{28,35} Therefore, the determination of APR can be viewed as a potentially valid measure of anaerobic capacity in handcyclists. In regard to upper body strength Nevin et al,²⁵ demonstrated that 8-weeks of concurrent strength and endurance training enhanced handcycling performance to a

greater degree than endurance training alone, suggesting that enhanced upper body strength may also be an important determinant of handcycling performance.

Given the current paucity of research relating to the impact of anaerobic threshold, anaerobic capacity and upper-body strength upon handcycling performance the aim of the present study was to build upon the existing literature and investigate how relative $\dot{V}O_{2peak}$, PO_{peak} , GME, power at a fixed blood lactate concentration of $4 \text{ mmol}\cdot\text{l}^{-1}$ (PO_4), $PO_{max,AO15}$, and measures of upper-body strength influence performance of an ecologically valid (field-based), 15-km TT in a group of trained H3/H4 male handcyclists. It was hypothesized that relative $\dot{V}O_{2peak}$, PO_{peak} , GME, PO_4 , $PO_{max,AO15}$, and relative upper-body strength would all demonstrate a significant correlation with 15-km TT velocity.

Methods

Participants

Thirteen male handcyclists with at least one year's recreational handcycling experience provided written informed consent to take part in this study. All participants were classified as either an H3 or H4 arm-powered handcyclist in accordance with current UCI Paracycling regulations.³³ Six participants were bi-lateral, above knee amputees (H4); one was a triple amputee (H3); and five were paraplegic with impairments corresponding to a spinal lesion between levels T1 to T10 (H3). In all aspects of physiological testing and TT performance, each participant used their own, customised handbike. Mean ($\pm SD$) participant characteristics were age 37 ± 11 yrs; body mass; 76.6 ± 10.1 kg; and 4-site skinfold summation 50.0 ± 7.2 mm. No medical conditions or upper-body musculoskeletal injuries were reported prior to the study. Finally, the study was conducted in accordance with the declaration of Helsinki with approval granted by the Research Ethics Committee of Buckinghamshire New University, High Wycombe, United Kingdom.

Design

This was a single-cohort, cross sectional research study to examine the influence of body mass, body composition, relative $\dot{V}O_{2peak}$, PO_{peak} , PO_4 , GME, $PO_{max,AO15}$, APR and upper-body strength on 15-km TT performance using thirteen experienced H3/H4 male handcyclists. Testing was completed over three consecutive days: 15-km TT (day 1), anthropometric assessment, graded exercise test (GTX), and 15-s all-out sprint test (day 2); and 1 repetition maximum (1RM) strength testing (day 3). A period of 24 hours separated testing sessions in order to limit the impact of fatigue. Before testing, participants were asked to abstain from strenuous exercise and refrain from consuming alcohol for at least 48 hours. Outdoor TT testing was conducted in dry and stable meteorological conditions ($19 \pm 2^\circ \text{C}$, $<10 \text{ km}\cdot\text{h}^{-1}$ wind speed) whilst, laboratory testing was performed indoors, under controlled environmental conditions (18°C , 50 – 60% relative humidity).

Individual 15-km Time Trial

In order to assess real world, ecologically valid handcycling performance, a 15-km individual TT was completed at a closed, cycling racing circuit (Odd Down, Bath, England). This location provided an undulating 1.5-km, smooth tarmac circuit with a total elevation loss and gain of 9 m per lap. Following two familiarisation laps, each participant was required to complete ten laps of the 1.5-km circuit as quickly as possible. Participants were monitored by means of a GPS receiver (Garmin Edge 1000, Garmin Ltd, USA), and data were used to establish TT performance in the form of mean velocity ($\text{km}\cdot\text{h}^{-1}$).

Anthropometry

Body mass was measured to the nearest 0.1 kg using a calibrated scale (Seca 714, Hamburg, Germany); whilst skinfold thicknesses were measured to the nearest mm using a pair of skinfold

callipers (accurate to 0.2 mm) from the Harpenden range of anthropometric instruments (Holtain, Ltd, UK). All anthropometric measurements including body mass and four-site skinfold thickness summation (chest, triceps, subscapular, and iliac crest), were performed in accordance with International Society for the Advancement of Kinanthropometry guidelines.¹⁷ Percentage body fat was not calculated as no validated four-site skinfold predication equations currently exists in the literature for disabled population groups with substantial body asymmetry as a result of amputation or lower body muscular atrophy due to spinal cord injury.¹⁵

Graded Exercise Test

For both the GTX and 15-s all-out sprint tests, each participant's bike was fitted to a standard, indoor cycling turbo trainer (Fluid 2, CycleOps, USA). Mechanical power output was measured using an instrumented front wheel hub (Powertap, G3, CycleOps, USA, 1.5% accuracy between 0 and 1999 W, sample frequency 0.2 Hz). The Powertap has been shown to be a reliable instrument (CV 0.9 – 2.9%) for the measurement of power whilst cycling⁷ and was calibrated prior to testing, in accordance with the manufacturer's instructions. Oxygen consumption ($\dot{V}O_2$), carbon dioxide production ($\dot{V}CO_2$), minute ventilation ($\dot{V}E$), and respiratory exchange ratio (RER) were continuously monitored using a calibrated, online gas analysis system (Oxycon Pro, Jeager, Warwick, Warwickshire, UK) whilst heart rate (HR) was logged using a commercially available receiver (Garmin Edge 1000, Garmin Ltd, USA).

Following a 10-min warm-up at a self-selected mechanical power output, participants were requested to start the test protocol at a work rate of 40 W with subsequent 20 W increments every 5-mins until the required mechanical power output could no longer be maintained, or until participants reached volitional exhaustion.^{2,27,31} $\dot{V}O_{2peak}$ and PO_{peak} were identified as the respective mechanical power output and peak oxygen consumption achieved during the last fully completed 30-s. Throughout the

test, participants were free to adjust their gear ratio and/or crank rate as needed in order to maintain the required mechanical power output. Every 5-mins and upon immediate completion of the test, participants were asked to indicate their global rating of perceived exertion (RPE) using a 6 to 20 Borg scale.⁸ All respiratory parameters were calculated for each breath and averaged at 1-min intervals at rest and every 30-s during each exercise stage.

At the end of each stage and at the point of volitional exhaustion, a small sample of capillary blood was collected from an earlobe to measure blood lactate concentration. These data were used to identify fixed blood lactate concentrations of 2 and 4 mmol·l⁻¹. Once collected, capillary blood samples were treated, analysed and disposed of immediately using a fully automated analyser (Biosen C-line, EKF Diagnostics, Barleben, Germany). GME was calculated as the ratio of external work produced to the amount of energy expended when a fixed blood lactate concentration of 2 mmol·l⁻¹ was reached. This metabolic parameter was selected as it represents a consistent, submaximal exercise intensity during which energy production is predominantly achieved via aerobic metabolic pathways. Metabolic energy expenditure was calculated from associated $\dot{V}O_2$ and RER data according to Garby and Astrup¹⁴ and expressed as a percentage value:

$$\text{Equation 1: GME} = ((\text{external work done} / \text{energy expenditure}) \times 100) (\%).$$

As an approximation of anaerobic threshold, power output corresponding to the onset of blood lactate accumulation (OBLA) at a fixed blood lactate concentration 4 mmol·l⁻¹ was also identified.

15-s All-Out Sprint Test

Following the GTX, participants were given a one-hour recovery period prior to completing a 15-s all-out sprint protocol to assess anaerobic performance. Participants were asked to complete a 10-

min warm up at a self-selected mechanical power output. Prior to commencement of the test protocol participants were requested to adopt their highest gear ratio (50/11). Once the participant acknowledged that they were ready, the test was initiated. Throughout the test protocol participants were verbally encouraged to exert maximum, physical effort with the greatest mechanical power output subsequently recorded. APR was established using the following formula:³⁵

$$\text{Equation 2: } \text{APR} = \text{PO}_{\text{max,AO15}} - \text{PO}_{\text{peak}}.$$

Upper-Body Strength Testing

In order to evaluate upper body strength, maximal and relative values of bench press and prone bench pull 1RM were determined. Strength testing was conducted on a specifically designed, IPC Para-powerlifting bench (Eleiko, Sweden) and a prone-pull bench (Pullum Sports, England) using a 20 kg Olympic barbell, 450 mm diameter barbell plates (25, 20, 15 and 10 kg), 200 mm diameter barbell plates (5, 2.5, 2.0, 1.5, 1.0 and 0.5 kg), two safety locks and two Velcro securing straps (Eleiko, Sweden).

Both bench press and prone bench pull 1RM testing was conducted in line with the protocols proposed by Haff and Triplett.¹⁶ Participants were instructed to perform a light warm-up with the bar only, performing 5 – 10 repetitions. Following a 1-min recovery period, a second set of 3 – 5 repetitions was performed with an estimated 60% 1RM load. After a 3-min recovery period another set of 2 – 3 repetitions, was performed with an estimated 80% 1RM load. Thereafter, an estimated 1RM load was selected and the participant asked to perform a single repetition. If successful, the participant was given a 3-min recovery period prior to performing a further 1RM attempt with an increased load.

Participants were allowed, to perform 3 – 5 more 1RM attempts with 3-min recovery between sets until their 1RM had been established within a precision of 1.0 kg.

Statistical Analyses

All data are reported as mean (\pm *SD*) with a level of significance for all statistical analyses set at $p < 0.05$. Statistical analyses were performed using SPSS Version 25.0 (SPSS Inc, Chicago, USA). Parameters were checked for normal distribution using the Shapiro-Wilk test with the Spearman's coefficient used in cases of violation. Pearson's product-moment correlation coefficients (r) were calculated to establish the relationships between 15-km TT velocity (dependent variable), body mass, 4-site skinfold summation, relative $\dot{V}O_{2peak}$, PO_{peak} , power-to-mass ratio ($W \cdot kg^{-1}$), PO_4 , GME, $PO_{max,AO15}$, APR, and maximal (kg) and relative ($kg \cdot kg^{-1}$ body mass) bench press and prone bench pull 1 RM (independent variables). Correlation coefficients were evaluated as follows >0.1 small, >0.3 moderate, >0.5 large, >0.7 very large, and >0.9 extremely large.¹⁷

Results

Mean (\pm *SD*) data are summarised in Table 1. Pearson product-moment correlation coefficients were calculated between 15-km TT velocity and all other anthropometric, GTX, 15-s all-out sprint and strength testing variables (Table 2). PO_4 ($r = .927$; $p < 0.01$) showed an extremely large correlation whilst relative $\dot{V}O_{2peak}$ ($r = .879$; $p < 0.01$), power-to-mass ratio ($r = .879$; $p < 0.01$), PO_{peak} ($r = .851$; $p < 0.01$), body mass ($r = -.783$; $p < 0.01$), GME ($r = .733$; $p < 0.01$), relative prone bench pull strength ($r = .770$; $p = 0.03$) relative bench press strength ($r = .703$, $p = 0.11$) and $PO_{max,AO15}$ ($r = .678$; $p = 0.15$) all demonstrated very large correlation with 15-km TT velocity. APR demonstrated a large correlation ($r = .548$; $p = 0.65$) whilst, body composition ($r = -.448$; $p = 0.14$), bench press 1RM ($r = .423$; $p = 0.17$) and prone bench pull 1RM ($r = .447$; $p = 0.14$) revealed only a moderate correlation with performance outcomes.

225

226 **##Insert Table 1 Here##**

227

228 **##Insert Table 2 Here##**

229

230 **##Insert Figure 1 Here##**

Accepted

Discussion

The aim of this study was to investigate the relationship between selected anthropometric, physiological, and upper body strength measures and 15-km handcycling time trial performance. The main findings based upon the data collected were that PO_4 , relative $\dot{V}O_{2peak}$, power-to-mass ratio, PO_{peak} , GME, relative upper body strength and $PO_{max,AO15}$ all demonstrated a significant correlation with handcycling performance (Figure 1).

Anthropometrics

In agreement with the findings of De Groot et al,⁹ the present study demonstrated that body mass displayed a considerable, negative association with 15-km TT velocity. When relative PO_{peak} was examined, another large and meaningful relationship was observed. These findings concur with those of Janssen et al,¹⁹ who demonstrated that power-to-mass ratio was a significant predictor of 10-km handcycling race performance. De Groot et al,⁹ reported that a lower waist circumference was a good predictor of handcycling performance, therefore, it can be inferred that a lower fat mass would result in improved 15-km TT performance. However, findings of the current study revealed that body composition, as assessed via 4-site skinfold summation exhibited only a moderate, negative relationship with performance. These findings were somewhat surprising as greater skinfold summation (*e.g.*, higher fat mass) would have been expected to have a larger negative impact upon 15-km TT velocity. Goosey-Tolfrey et al,¹⁵ suggested that skinfold estimations may not be a valid assessment tool of body composition in disabled population groups and the findings of the present study support this position. Another point to note is that the combined rider-bike mass is also likely to be important and negatively linked to TT velocity. Thus, where feasible, it is recommended that the combined mass of a handcyclist and their bike be minimised in an attempt to optimise TT performance.

256 **Aerobic Performance**

257 Relative $\dot{V}O_{2\text{peak}}$ and, to a slightly lesser extent PO_{peak} and GME demonstrated a very large correlation
 258 with 15-km TT performance. These finding are in agreement with previous studies that have shown
 259 relative $\dot{V}O_{2\text{peak}}$, and PO_{peak} to be significant determinants of handcycling performance during
 260 TT,^{13,18,19} ultra-endurance,^{1,2} and mountain climbing,⁹ events. GME has also been demonstrated to be
 261 a significant determinant of handcycling performance.¹⁹ Participants in the present study achieved a
 262 mean GME value of $13.4 \pm 2.7\%$ which is similar to a previous reports of $11.5 \pm 0.8\%$,¹⁹ $14.1 \pm$
 263 2.0% ,²¹ and $13.5 \pm 1.4\%$ ²⁵ for trained handcyclists. GME is of particular importance to handcyclists
 264 as improvements in efficiency will likely translate to a reduction in relative workload at a given
 265 mechanical power output. Theoretically, this should enable a rider to produce a higher mechanical
 266 power output for an equivalent amount of energy expended (*e.g.*, improved performance) or
 267 alternatively result in a longer time to exhaustion at a given rate of work (*e.g.*, improved endurance
 268 capacity), with both scenarios holding the potential to enhance an athlete's performance.

270 **Anaerobic Threshold**

271 As an approximation of anaerobic threshold, mechanical power output corresponding to OBLA at a
 272 fixed blood lactate concentration of $4 \text{ mmol}\cdot\text{l}^{-1}$ (PO_4) was utilised. Power at this threshold
 273 demonstrated an extremely large relationship with 15-km TT velocity. These finding are in agreement
 274 with findings of several other studies, demonstrating a strong relationship between PO_4 and
 275 performance in both handcyclists^{2,23} and able-bodied road cyclists.^{10,11} OBLA can be defined as the
 276 point at which blood lactate concentration increases exponentially, with the rate of production
 277 outstripping removal and leading to acute metabolic acidosis. The validity of using OBLA at a fixed
 278 blood lactate concentration of $4 \text{ mmol}\cdot\text{l}^{-1}$ to identify the shift from predominantly aerobic energy
 279 turnover to anaerobic energy production has been questioned previously.^{10,11,30} Indeed, the threshold
 280 at which anaerobic metabolism starts to predominate is strongly influenced by individual lactate

281 kinetics, which varies depending upon the activity and volume of skeletal muscle mass.²⁶ Therefore,
282 OBLA at a fixed blood lactate concentration of 4 mmol·l⁻¹ may either over- or underestimate
283 anaerobic threshold depending upon the individual athlete, exercise modality and intensity.
284 Nevertheless, findings of the present study demonstrated that anaerobic threshold, as estimated by
285 PO₄ to be a very strong indicator of 15-km handcycling TT performance.

286

287 **Anaerobic Performance**

288 Anaerobic performance assessed by the maximum power output generated during a 15-s all-out sprint
289 test produced a large correlation with 15-km TT performance. In order to assess anaerobic capacity,
290 the difference between PO_{max,AO15} and PO_{peak} commonly known as APR was also investigated. The
291 concept of APR has been suggested to be a valid measure of anaerobic capacity in able-bodied road
292 cyclists^{28,35} and findings of the current study showed large correlation between handcycling APR and
293 15-km TT velocity. Given the nature of an individual TT, it can be reasoned that riders in the present
294 study maintained a velocity at, or just below their anaerobic threshold and were unlikely to have
295 utilised their APR to a great extent. Despite this, it can be strongly argued that in the context of a road
296 race competitive handcyclists require a high APR to repeatedly generate and recover from the
297 production of high power outputs over short periods of time. This will allow a rider to either close a
298 gap on an opponent, break away from other riders, or win in a sprint finish.

299

300 **Maximal and Relative Upper Body Strength**

301 Maximal and relative upper body strength in the present study was determined by assessing bench
302 press and prone bench pull 1RM. These exercises were chosen as they closely mimic the
303 synchronistic, horizontal push/pull force production movement pattern observed during
304 handcycling.²⁵ Bench press and prone bench pull 1RM demonstrated only a moderate relationship
305 with 15-km TT velocity. However, when relative upper body strength was examined, both bench

press and prone bench pull displayed a significant influence upon 15-km TT velocity, further confirming the relative importance of body mass to handcycling performance.

Relative strength is the product of one's ability to generate considerable maximal forces relative to one's body mass. Therefore, it can be inferred that greater maximal upper body strength, at a given body mass or in combination with a reduction in non-functional body mass (*e.g.*, adipose tissue) may result in an improvement in an individual's handcycling performance capabilities. Interestingly, relative upper body pulling strength demonstrated a larger correlation with 15-km TT velocity than relative pushing strength. This is in agreement with previously published studies which have demonstrated that a greater proportion of the work generated during the propulsion cycle of handcycling occurs during the pull phase with an increase in pulling torque and concomitant decrease in pushing torque observed at progressively higher power outputs.^{5,27,34} Therefore, based upon these observations it can be inferred that greater relative upper body pulling strength may enhance handcycling performance. Strength training should, therefore, form a central component of any successful handcyclists training programme.

Limitations

While these findings provide a novel insight into the determinants of TT performance in trained H3/H4 handcyclists, it must be noted that there are several limitations associated with the study and its design. Firstly, as is usually the case in this area of research, sample size was small and the participant group relatively heterogeneous in terms of age, performance level, and disability which resulted in considerable variance within the group. The small sample size also meant that it was not possible to conduct a multiple regression analysis in order to develop a accurate handcycling performance model. A further limitation was that the 15-km TT was a self-paced time trail, which was conducted in variable climactic conditions. Such an approach represents a less controlled

environment compared to sterile laboratory conditions; however, this approach does add a degree of ecological validity as it more closely resembles real-world handcycling race conditions. Additionally, whilst the Powertap power measuring device has been shown to be a reliable instrument for the measurement of mechanical power output whilst cycling, its sampling frequency (0.2 Hz) is relatively low for the purpose of measuring $PO_{\max, AO15}$. Thus, it is recommended that future studies use a more sensitive instrument for the measurement of mechanical power output such as the Cyclus 2 ergometer (8 Hz, RBM Electronic automation GmbH, Leipzig, Germany). However, it could also be argued that, from a consistency of power measurement perspective and to facilitate meaningful comparison, the same measuring device should be used in a laboratory and field-based environment if possible. Finally, as with all criterion validity studies exploring links between independent variables and, in this case, a (dependent) performance outcome measure, correlational analyses do not necessarily confirm causation, thus additional studies should be conducted to provide further, confirmatory evidence.

Practical Applications

In order to better optimise handcycling performance capability, it is recommended that an emphasis be placed upon the development and frequent monitoring of the following parameters: PO_4 , relative $\dot{V}O_{2peak}$, power-to-mass ratio, PO_{peak} , GME, relative upper body strength and $PO_{\max, AO15}$. Particular attention should be placed upon the development of upper body pulling strength in order to enhance the force production during the pulling phase of the handcycling propulsion cycle. Finally, whilst not confirmed by the findings of this study, riders should aim to reduce their overall body fat summation in order to further improve handcycling performance capabilities. Linked to this point, it follows that it would also be prudent for a competitive handcycle to be as lightweight as possible

Conclusion

In conclusion, the findings of the present study indicate that PO_4 , relative $\dot{V}O_{2peak}$, power-to-mass ratio, PO_{peak} , GME, relative upper body strength, $PO_{max,AO15}$ and APR all have significant impact upon 15-km TT velocity in H3/H4 handcyclists. To the best of the authors' knowledge, this is the first study to investigate the combined impact of anaerobic threshold, anaerobic capacity and upper-body strength upon real-world handcycling TT performance. Based upon our findings it is recommended that future research associated with establishing determinants of handcycling performance use a larger, more homogenous, group of competitive, preferably elite handcyclists. If sufficient data could be collected it would be possible to construct an accurate performance model using multiple regression analysis.

Declaration of Interest

There is no conflict of interest to declare for the present article.

References

1. Abel T, Schneider S, Platen P, & Struder HK. Performance diagnostics in handcycling during competition. *Spinal Cord*. 2006; 44: 211 – 216.
2. Abel T, Burkett B, Schneider S, et al. The exercise profile of an ultra-long handcycling race: The Stykkeproven experience. *Spinal Cord*. 2010; 48: 894 – 898.
3. Abel T, Vanlandewijck Y, & Verellen J. Handcycling, In *Wheelchair Sport*. Champaign, IL: Human Kinetics; 2010.
4. Abel T, Burkett B, Thees B, et al. Effect of three different grip angles on physiological parameters during 2 laboratory handcycling tests in able-bodied participants. *Front Physio*. 2015; 23(6): 331.

- 380 5. Arnet U, Van Drongelen S, Van Der Woude LHV, et al. Shoulder load during handcycling at
381 different incline and speed positions. *Clin Biomech.* 2012; 27(1): 1 – 6.
- 382 6. Arnet U, Van Drongelen S, Schlussel, et al. The effect of crank position and backrest
383 inclination on shoulder load and mechanical efficiency during handcycling. *Scand J Med Sci*
384 *Sport.* 2014; 24: 386 – 394.
- 385 7. Bertucci W, Duc S, Villerius V, et al. Validity and reliability of the Powertap mobile cycling
386 power meter when compared with the SRM device. *Int J Sports Med.* 2015; 26(10): 868 –
387 873.
- 388 8. Borg, GA. Psychophysical based of perceived exertion. *Med Sci Sports Exerc.* 1982; 14: 377
389 – 388.
- 390 9. De Groot S, Postma K, Van Vliet L, et al. Mountain time trial time in handcycling: exercise
391 intensity and predictors of race time in people with spinal cord injury. *Spinal Cord.* 2014; 52:
392 455 – 461.
- 393 10. Faria EW, Parker DL,& Faria IE. The science of cycling physiology and training – Part 1.
394 *Sports Med.* 2005; 35(4): 313 – 337.
- 395 11. Faria, EW, Parker DL, & Faria IE. The science of cycling factors affecting performance – Part
396 2. *Sports Med.* 2005; 35(4): 313 – 337.
- 397 12. Faupin A, Gorce P, Watelain E, et al. A biomechanical analysis of handcycling: A case study.
398 *J Appl Biomech.* 2010; 2: 240 – 245.
- 399 13. Fischer G, Ardigo L, & Figueiredo P. Physiological performance determinants of a 22-km
400 handbiking time trail. *Int J Sports Physiol Perf.* 2016; 10: 965 – 971.
- 401 14. Garby L, & Astrup A. The relationship between the respiratory quotient and the equivalent of
402 oxygen during simultaneous glucose and lipid oxidation and lipogenesis. *Acta Physiol Scandi.*
403 1987; 129(3): 443 – 444.

- 404 15. Goosey-Tolfrey V, Keil M, Brooke-Wavell K, & De Groot S. A comparison of methods for
405 the estimation of body composition in highly trained wheelchair games players. *Int J Sports*
406 *Med.* 2016. 37(10): 799 – 806.
- 407 16. Haff GG, & Triplett NT. *Essentials of Strength Training and Conditioning* (4th Ed).
408 Champaign, Il: Human Kinetics. 2016.
- 409 17. Hopkins W, Marshall S, Batterham A, et al. Progressive statistics for studies in sports
410 medicine and exercise science. *Med Sci Sports Exerc.* 2009; 14: 3 – 12.
- 411 18. International Society for the Advancement of Kinanthropometry. *International Standards for*
412 *Anthropometric Assessment.* 2001.
- 413 19. Janssen TWJ, Dallmeijer AJ, & Van der Woude LHV. Physical capacity and race
414 performance of handcycle users. *J Rehabil Res Develop.* 2001; 38(1): 33 – 40.
- 415 20. Litzenberger S, Mally, F, & Sabo, A. Biomechanics of elite recumbent handcycling. A case
416 study. *Sports Eng.* 2016; 19(3), 201 – 211.
- 417 21. Lovell D, Sheilds D, Beck B, Cuneo R, et al. The aerobic performance of trained and
418 untrained handcyclists with spina cord injury. *Euro J Appl Physiol.* 2012; 112(9): 3431 –
419 3437.
- 420 22. Mannion P, Toparlar Y. Clifford E, Hajdukiewicz M, Andrainne T, & Blocken B. The impact
421 of arm-crank position on the drag of a Paralympic hand-cyclist. *Comp Methods in Biomech*
422 *and Biomech Engr.* 2019; 22(4): 386 – 395.
- 423 23. Perrett, C. Elite-adapted wheelchair sports performance: a systematic review. *Disabil Rehabil.*
424 2017; 39(2): 164 – 172.
- 425 24. Mason BS, Van der Woude LHV, & Goosey-Tolfrey VL. The ergonomics of the wheelchair
426 configuration for optimal performance in the wheelchair court sports. *Sports Med.* 2013;
427 43(1): 164 – 172.

- 428 25. Nevin JP, Smith P, Waldron M, et al. Efficacy of an 8-week concurrent strength and
429 endurance training programme on hand cycling performance. *J Strength Cond Res.* 2018;
430 32(7): 1861 – 1868.
- 431 26. Quittman OJ, Abel T, Zeller S, et al. Lactate kinetics in handcycling under various exercise
432 modalities and their relationship to performance measures of able-bodied participants. *Eur J*
433 *Appl Physiol.* 2018; 118(7): 1493 – 1505.
- 434 27. Quittman OJ, Meskemper, J, Abel, T et al. Kinematics and kinetics of handcycling propulsion
435 at increasing workloads in able-bodied subjects. *Sports Eng.* 2018; 21(4): 283 – 294.
- 436 28. Sanders D, & Heijboer, M. The anaerobic power reserve and its applicability in professional
437 road cycling. *J Sports Sci.* 2019; 37(6): 621 – 629.
- 438 29. Smith P, Chapman M, Hazelhurst K, et al. The influence of crank configuration on muscle
439 activity and torque production during arm crank ergometry. *J Electro Kinesio.* 2008; 18(4): 598
440 – 605.
- 441 30. Stangier C, Abel T, Zeller S, Quittman OJ, Perret C, & Struder HK. Comparison of different
442 blood lactate threshold concepts for constant load performance in spinal cord injury
443 handcyclists. *Front Physio.* 2019; 10 (1054): doi: 10.3389/fphys.2019.01054.
- 444 31. Stone B, Mason BS, Warner MB, et al. Shoulder and thorax kinematics contribute to increased
445 power output of competitive handcyclists. *Scand J Med Sci Sport.* 2019; 29(6): 843 - 853.
- 446 32. Stone B. Mason BS, Warner, MB & Goosey-Tolfrey VL. Horizontal crank position affects
447 economy and upper limb kinematics of recumbent handcyclists. *Med Sci Sports Exerc.* 2019;
448 51(11), 2265 – 2273.
- 449 33. Union Cycliste Internationale, *Cycling Regulations*, Part 16 Para-Cycling. 2019.
- 450 34. Vegter RJK, Mason, BS, Sporrel B, Stone B, Van Der Woude, LHV & Goosey-Tolfrey VL.
451 Crank fore-aft position alters the distribution of work over the push and pull phase during

synchronous recumbent handcycling in able-bodied participants. *PloS one*. 2019; 14(8): doi:
10.1371/journal.pone.0220943.

35. Weyand, PG, Line JE, & Bundle MW. Sprint performance duration relationships are set by the
fractional duration of external force application. *Am J Physio*. 2006; 290(3): 758 – 765.

Accepted

Table 1. Mean (\pm SD) values of participant testing data

15 km Time Trial Velocity ($\text{km}\cdot\text{h}^{-1}$)	28.6 ± 6.3
Peak Heart Rate (bpm)	174 ± 12
Peak Blood Lactate ($\text{mmol}\cdot\text{l}^{-1}$)	11.9 ± 1.8
$\dot{V}\text{O}_{2\text{peak}}$ ($\text{l}\cdot\text{min}^{-1}$)	2.8 ± 0.5
Relative $\dot{V}\text{O}_{2\text{peak}}$ ($\text{ml}\cdot\text{kg}^{-1}\text{min}^{-1}$)	36.8 ± 10
PO_{peak} (W)	160 ± 26.7
Relative PO_{peak} ($\text{W}\cdot\text{kg}^{-1}$)	2.2 ± 0.7
GME (%)	13.4 ± 2.7
PO_4 (W)	119 ± 26
$\text{PO}_{\text{max,AO15}}$ (W)	547 ± 120
APR (W)	387 ± 107
Bench Press 1RM (kg)	90.2 ± 16.7
Relative Bench Press Strength ($\text{kg}\cdot\text{kg}^{-1}$ body mass)	1.2 ± 0.3
Prone Bench Pull 1RM (kg)	77.8 ± 13.2
Relative Prone Bench Pull Strength ($\text{kg}\cdot\text{kg}^{-1}$ body mass)	1.0 ± 0.3

Table 2. Correlations among parameters of 15-km TT performance

	Velocity (km·h ⁻¹)	Body Mass (kg)	4-Site Skin Fold Summation (mm)	Relative VO _{2peak} (ml·kg ⁻¹ ·min ⁻¹)	PO _{peak} (W)	Power-to- Mass Ratio (W·kg ⁻¹)	GME (%)	PO ₄ (W)	PO _{max,AO15} (W)	APR (W)	Bench Press 1RM (kg)	Relative Bench Press Strength (kg·kg ⁻¹ body mass)	Prone Bench Pull 1RM (kg)	Relative Prone Bench Pull Strength (kg·kg ⁻¹ body mass)
Velocity (km·h ⁻¹)	-													
Body Mass (kg)	-.783**	-												
4-Site Skin Fold Summation (mm)	-.448	.151	-											
Relative VO _{2peak} (ml·kg ⁻¹ ·min ⁻¹)	.879**	-.835*	-.510*	-										
PO _{peak} (W)	.851**	-.825**	-.494	.774**	-									
Power-to-Mass Ratio (W·kg ⁻¹)	.879**	-.921**	-.391	.831**	.964**	-								
GME (%)	.733**	-.821**	-.528	.853**	.717**	.798**	-							
PO ₄ (W)	.927**	-.668*	-.637	.861**	.842**	.827**	.709**	-						
PO _{max,AO15} (W)	.678*	-.571	-.383	.775**	.572	.563	.641*	.595*	-					
APR (W)	.548	-.435	-.306	.676*	.392	.392	.540	.457	.979**	-				
Bench Press 1RM (kg)	.423	-.505	-.271	.603*	.310	.379	.686*	.346	.684*	.690*	-			
Relative Bench Press Strength (kg·kg ⁻¹ body mass)	.703*	-.814**	-.313	.822**	.647*	.735**	.871**	.615*	.734**	.662*	.899**	-		
Prone Bench Pull 1RM (kg)	.447	-.363	-.331	.506	.275	.285	.498	.358	.566	.657	.865**	.728**	-	
Relative Prone Bench Pull Strength (kg·kg ⁻¹ body mass)	.770**	-.793**	-.356	.819**	.671**	.734**	.811**	.661*	.701*	.619*	.852**	.949**	.843**	-

**Correlation is significant at the <0.01 level (2-tailed)

* Correlation is significant at the < 0.05 level (2-tailed)




Fig 1. Correlation plots between 15-km TT velocity and parameters of exercise testing